



New erymid lobsters from the Nusplingen and Usseltal formations (Upper Jurassic) of southwest Germany

Novi jastogi iz družine Erymidae iz formacij Nusplingen in Usseltal (zgornja jura) iz jugozahodne Nemčije

Günter SCHWEIGERT¹ & Jürgen HÄRER²

¹Staatliches Museum für Naturkunde, Rosenstein 1, 70191 Stuttgart, Germany; guenter.schweigert@smns-bw.de

²Roennebergstraße 5, 12161 Berlin, Germany; juergen.haerer@gmail.com

Prejeto / Received 28. 10. 2019; Sprejeto / Accepted 2. 4. 2020; Objavljeno na spletu / Published online 5. 7. 2020

Key words: Decapoda, Erymidae, Kimmeridgian, Tithonian, Plattenkalk, taphonomy

Ključne besede: dekapodni raki, Erymidae, kimmeridgij, tithonij, litografski apnenec, tafonomija

Abstract

Two new species of Late Jurassic erymid lobsters, *Stenodactylina devillezi* sp. nov. and *Stenodactylina geigerae* sp. nov., are described on the basis of isolated, but well-preserved chelipeds from the upper Kimmeridgian of Swabia and the lower Tithonian of Bavaria, respectively. The incomplete nature of the material indicates that these lobsters were not autochthonous elements of the Plattenkalk lagoons, but resulted from predation in nearby shallow-water settings.

Izvilleček

V prispevku predstavljamo dve novi vrsti zgornjejurskih jastogov iz družine Erymidae, to sta *Stenodactylina devillezi* sp. nov. in *Stenodactylina geigerae* sp. nov. Novi vrsti sta opisani na podlagi izoliranih, vendar dobro ohranjenih škarnikov iz zgornjega kimmeridgija Švabske in spodnjega tithonija Bavarske. Slabša ohranjenost primerkov kaže, da tovrstni jastogi niso bili avtohtoni prebivalci zgornje jurskih lagun, ampak so verjetno poseljevali bližnja plitvodna okolja.

Introduction

The Upper Jurassic Solnhofen-type lithographic limestones (“plattenkalks”) of southwest Germany are renowned for their exceptionally preserved fossils, making them one of the classic examples of conservation Fossilagerstätten (Seilacher et al., 1985). In addition to iconic vertebrate fossils such as the ancient bird *Archaeo-ptyx* von Meyer, 1861, hundreds of taxa have been recorded over the centuries (e.g., Leich, 1968; Barthel et al., 1990; Frickhinger, 1994, 1999; Arratia et al., 2015). However, it should be noted that fossils labelled “Solnhofen” come from various deposits of different lithology and age (Schweigert, 2007, 2015a). Decapod crustaceans from these lithographic limestones, and erymid lobsters in particular, have been described from the early 19th century onwards (e.g.,

Desmarest, 1817, 1822; von Schlotheim, 1822; zu Münster, 1839; Oppel, 1861, 1862). More recently, Beurlen (1928) and Förster (1966) studied Jurassic erymids. However, some of their systematic assignments remained questionable. Taxa based on material from shallow-water deposits consist either of carapace remains or of isolated chelae. More complete specimens, with the carapace and corresponding chelae preserved, are only found when conditions were favourable or within concretions (e.g., Hyžný et al., 2015). At generic level, erymid lobsters are distinguished mainly by their characteristic carapace groove pattern (Devillez & Charbonnier, 2017, 2019, 2020; Devillez et al., 2018), while chela shape is less characteristic, at least in *Eryma* s. str. In the latter, there is one clade that has chelae with relatively short fingers; this group is distinguished

from another with strikingly longer, straight fingers, although the groove pattern of the carapace is the same. As a result, some species of *Eryma* of the second clade, previously included in a distinct genus, *Erymastacus* Beurlen, 1928 (see Hyžný et al., 2015), have later been considered to belong to *Eryma* s. str. (Devillez & Charbonnier, 2017, 2019, 2020). However, there are further taxa that have long and curved fingers, but a different groove pattern on their carapace. These are now included in the genus *Stenodactylina* Beurlen, 1928. Remains of *Stenodactylina* are easily recognised on the basis of their chelae even if such occur isolated (Devillez & Charbonnier, 2019a, 2019b). Most Late Jurassic remains of *Stenodactylina* represent chelae or chelipeds from coral-bearing limestones or other shallow-water lithologies (e.g., Étallon, 1859, 1861; Krause, 1891; Bachmayer, 1959). Here we describe the first examples of *Stenodactylina* from Upper Jurassic lithographic limestones.

Geological settings

Two of the specimens of *Stenodactylina* described here come from the upper Kimmeridgian Nusplingen Plattenkalk, also known as Nusplingen Lithographic Limestone. The site of prove-

nance is located in the western part of the Swabian Alb (Fig. 1). The Nusplingen Plattenkalk was deposited in a c. 80–100 metres deep lagoon surrounded by shallower areas and small islands (Stevens et al., 2014). Known since the mid-19th century, it has meanwhile provided more than 400 fossil taxa, among them pterosaurs, marine crocodiles, sharks and numerous other fishes, but also squid-like teuthoids and decapod crustaceans (Dietl & Schweigert, 2011). Ammonites have allowed to date it as late *Beckeri* Zone, *Ulmense* Subzone, *hoelderi* Biohorizon (Schweigert, 2007, 2015a). Especially common are large-sized penaeid prawns (Schweigert, 2001b, 2017; Odin et al., 2019), but polychelid, glypheid and erymid lobsters have been recorded as well (Fraas, 1855; Oppel, 1861, 1862; Schweigert & Dietl, 1999; Schweigert et al., 2000; Schweigert, 2001a; Charbonnier et al., 2013; Audo et al., 2014). Erymid material from Nusplingen is often incomplete due to predation or decay; however, several valid taxa have been described on the basis of such incomplete specimens (Oppel, 1861, 1862; Schweigert et al., 2000). Recently, this material has been re-studied within the context of a comprehensive review of all Late Jurassic erymid lobsters (Devillez & Charbonnier, 2020).




Fig. 1. Localities that yielded the material of *Stenodactylina* (asterisks) and additional Upper Jurassic Plattenkalk localities in southwest Germany (modified from Fürsich et al., 2007).

A third specimen of *Stenodactylina* comes from Upper Jurassic plattenkalks exposed at the northern hillside of the River Danube, southwest of the village of Marxheim (Fig. 1). These plattenkalks had not been studied previously for their fossil content. Fesefeldt (1962) mapped the area around Marxheim and described several sections and outcrops. From his descriptions the plattenkalks in question match the so-called “Spindeltal-Schiefer”, an informal lithological unit which is included in the lower Tithonian Usseltal Formation (Zeiss, 1977; Niebuhr & Pürner, 2014). For a determination of the age of these plattenkalks, several ammonite remains were sampled from the scree; these include *Subplanitoides spin-delense* Zeiss, 1968 (Fig. 2), *Subplanitoides* sp., “*Torquatisphinctes*” *regularis* Zeiss, 1968 and *Usseliceras* sp. This association clearly indicates the *franconicum* Biohorizon of the lower Tithonian *Mucronatum* Zone (see Schweigert, 2015a). The Submediterranean *Mucronatum* Zone corresponds approximately to the Tethyan *Darwini* Zone (Scherzinger & Schweigert, 2003). Assuming the duration of a biohorizon to have been around 165 ka (Schweigert, 2006), the Tithonian type horizon of the specimen from Marxheim is around 1.15 myr younger than the Kimmeridgian Nusplingen site.

Methods

The specimens studied were carefully prepared mechanically with needles and scalpels using a binocular with 50 × magnification. Photographs were taken with digital cameras under normal

or ultraviolet illumination. Ultraviolet illumination is often used to enhance the contrast between phosphatic fossils and the surrounding rock matrix (e.g., Haug et al., 2009; Tischlinger, 2015). The photographs were finally mounted as illustrations using Adobe Photoshop version CS5.1.

Systematic palaeontology

Class Malacostraca Latreille, 1802

Order Decapoda Latreille, 1802

Superfamily Erymoidea Van Straelen, 1925

Family Erymididae Van Straelen, 1925

Genus *Stenodactylina* Beurlen, 1928

Included species: *Stenodactylina armata* (Secretan, 1964), *S. australis* (Secretan, 1964), *S. burgundiaca* (Crônier & Courville, 2004), *S. delphinensis* (Moret, 1946), *S. devillezi* sp. nov. (herein), *S. falsani* (Dumortier, 1867), *S. geiger-ae* sp. nov. (herein), *S. guisei* (Wright, 1881), *S. insignis* (Oppel, 1862), *S. lagardettei* (Hyžný, Schlögl, Charbonnier, Schweigert, Rulleau & Gouttenoire, 2015), *S. liasina* Beurlen, 1928 (type species), *S. rogerfurzei* Schweigert, 2013, *S. spinosa* (Étallon, 1861), *S. strambergensis* (Bachmayer, 1959), *S. triglypta* (Stenzel, 1945) and *S. walkerae* (Feldmann & Haggart, 2007).

Stenodactylina devillezi sp. nov.

2003 *Erymastacus* sp. nov. – Schweigert & Garassino, fig. 2B.

2020 [fragmentary] *Stenodactylina*. – Devillez & Charbonnier, in press.




Fig. 2. The ammonite *Subplanitoides spindelense* Zeiss, 1968 from the lower Tithonian, *Mucronatum* Zone (= *Darwini* Zone), ‘Spindeltalschiefer’ of Usseltal Formation, Marxheim (leg. K. Geiger). Photograph: K. Geiger. Scale bar equals 20 mm.



Fig. 3. *Stenodactylina devillezi* sp. nov., A, holotype, SMNS 64872; B, paratype, SMNS 70506. Upper Kimmeridgian, *Beckeri* Zone, *Ulmense* Subzone, Nusplingen Formation, Nusplingen Quarry, Westerberg hill west of Nusplingen, southwest Germany. Photographs: G. Schweigert. Scale bars equal 10 mm.

Holotype: SMNS 64872, from the Nusplingen Formation (upper Kimmeridgian, *Beckeri* Zone) of Nusplingen, Baden-Württemberg, southwest Germany (Fig. 1).

Paratype: SMNS 70506, from same locality and biohorizon as the holotype.

Etymology: Named after Julien Devillez (Paris), who revised all previously described erymid taxa from the Jurassic and Cretaceous.

Type locality and horizon: Nusplingen Quarry, west of Nusplingen, southwest Swabian Alb (Fig. 1); Nusplingen Formation (upper Kimmeridgian, *Beckeri* Zone, *Ulmense* Subzone; see Schweigert, 2007).

Diagnosis: Species of *Stenodactylina* with a P1 chela that is characterised by a slender subrectangular manus, a completely toothless dactylus

and an index with seven teeth in the proximal half and a sinuosity in the distal part.

Description: The holotype is a relatively large left cheliped consisting of a well-preserved chela with manus and remains of the carpus. Manus subrectangular, length 33 mm, width 17 mm. Manus and carpus covered with fine, randomly scattered pustules; fingers lacking pustules, only with a few setal pits. Dactylus length 34 mm; index and dactylus terminally curved inwards. Occlusal surface of dactylus toothless; index bearing seven teeth in its proximal part, the strongest one is the third as counted from the distal side, with a longer distance between the first and the second. Distal half of index with a sinuosity.

The paratype is a right cheliped consisting of a manus with chela, the carpus and merus. Manus




Fig. 4. *Stenodactylina geigerae* sp. nov., holotype, SMNS 70507. A: photographed under normal illumination; B, photographed under ultraviolet illumination. Lower Tithonian, *Mucronatum* Zone (= *Darwini* Zone), 'Spindeltalschiefer' of Usseltal Formation, Marxheim, southwest Germany. Photographs: J. Härer. Scale bars equal 10 mm.

subrectangular, length 22 mm, width 10.5 mm. Carpus 10.5 mm long; merus poorly preserved, 26 mm long. Ornamentation of manus and carpus identical to that in the holotype. Dactylus length 26 mm, with some setal pits, otherwise smooth, distally curved, slightly dislocated during burial. Index mostly covered by dactylus, distal half broken off.

Remarks: The slight differences in the fingers/manus length ratio between the left and right che-liped (although observed in different individuals) point to heterochely in *S. devillezi* sp. nov., which is well known in other species of *Stenodactylina* (see Hyžný et al., 2015). A very fragmentary chela from Nusplingen that shows long and slender fingers with strong and widely spaced teeth,

originally recorded in open nomenclature (Schweigert et al., 2000: pl. 5, fig. 4, as “Erymidæ gen. et. sp. indet.”), possibly belongs to *Enoploclytia* M’Coy, 1849, another erymid genus recently recorded from the Upper Jurassic (Devillez et al., 2018). The rather atypical, slender appearance of the propodus is caused by incomplete preservation. No further specimen has been found so far.

***Stenodactylina geigeræ* sp. nov.**

Holotype: SMNS 70507, from the Usseltal Formation (lower Tithonian, *Mucronatum* Zone) of Marxheim, Bavaria, southwest Germany (Fig. 1).

Etymology: Named after Katharina Geiger (Munich), who found and kindly donated the specimen.

Type locality and horizon: Northern hillside of the River Danube, southwest of Marxheim, Bavaria, Germany (Fig. 1), Usseltal Formation, lower Tithonian, *Mucronatum* Zone (= Submediterranean equivalent of Tethyan *Darwini* Zone).

Diagnosis: Species of *Stenodactylina* with a P1 chela that is characterised by only two teeth in the proximal third of the dactylus and index.

Description: The holotype is a left cheliped with the manus, carpus and merus preserved. Several fine cracks are detectable resulting from compaction of the hollow fossil. Manus subrectangular, 19 mm long, 9.5 mm wide. Merus, carpus and manus covered with randomly scattered small pustules; coarser tubercles occur only along the inner margins of the merus and carpus. Merus length 23 mm, with a few spiny tubercles along the articulation towards the carpus. Carpus length 12 mm, distally bordered by a smooth seam. Fingers c. 28 mm long, lacking any pustules, only bearing a few tiny setal pits. Occlusal surfaces of dactylus and index each with two prominent teeth in the most proximal parts, otherwise smooth. Strongest tooth is the first one, counted from the proximal side of each finger.

Discussion

The preservation of isolated chelipeds of *Stenodactylina* in lithographic limestones probably results from predation activity in nearby shallower environments. The lobsters themselves usually did not inhabit the hostile sea floor of the lagoons. The most common way for erymid lobsters and most other decapod crustaceans to become fossilised complete in Solnhofen-type plattenkalks is via exuviae (Schweigert & Garassino, 2003). The bulk of exuviae, however, represent juvenile specimens, whereas larval stages are rare

as well due to their poor sclerotization (Haug et al., 2011). Juvenile stages of erymid lobsters are not easy to differentiate and much more material is needed to reconstruct their ontogenies.

Acknowledgements

Katharina Geiger (Munich) is thanked for donating one the specimens described here and for providing useful information on the locality. Martin Kapitzke and Markus Rieter (both SMNS) skilfully prepared the two specimens from Nusplingen. We also thank the volunteers, namely Gerd Dietl (Stuttgart), Rolf Hugger (Albstadt-Onstmettingen), August Ilg (Düsseldorf) and Burkhard Russ (Nusplingen), who helped excavate the Nusplingen Plattenkalk during the last 25 years and who recovered numerous rare specimens such as the present ones. The journal reviewers, Sylvain Charbonnier (Paris) and Julien Devillez (Paris), provided helpful comments for improvement of the manuscript.

References

- Arratia, G., Schultze, H.-P., Tischlinger, H. & Viohl, G. 2015: Solnhofen - Ein Fenster in die Jurazeit. Friedrich Pfeil, München: 620 p.
- Audo, D., Schweigert, G., Haug, J.T., Haug, C., Saint Martin, J.-P. & Charbonnier, S. 2014: Diversity and palaeoecology of the enigmatic *Knebelia* (Eucrustacea, Decapoda, Eryonidae) from the Upper Jurassic plattenkalks of southern Germany. *Palaeontology*, 57/2: 397-416. <https://doi.org/10.1111/pala.12071>
- Bachmayer, F. 1959: Neue Crustaceen aus dem Jura von Stramberg (ČSR). *Sitzungsberichte der Österreichischen Akademie der Wissenschaften, mathematisch-naturwissenschaftliche Klasse, Abteilung I*, 168: 937-944.
- Barthel, K.W., Swinburne, N.H.M. & Conway Morris, S. 1990: Solnhofen: a study in Mesozoic palaeontology. Cambridge University Press, Cambridge: IX+236 p.
- Beurlen, K. 1928: Die Decapoden des Schwäbischen Jura mit Ausnahme der aus den oberjurassischen Plattenkalken stammenden. *Palaeontographica*, 70: 115-278.
- Charbonnier, S., Garassino, A., Schweigert, G. & Simpson, M. 2013: A worldwide review of fossil and extant glypheid and litogastrid lobsters (Crustacea, Decapoda, Glypheoidea). *Mémoires du Muséum national d'Histoire naturelle Paris*, 205: 304 p.

- Crônier, C. & Courville, P. 2004: A rich and highly endemic decapod crustacean fauna from the Middle Jurassic of north-east France. *Palaeontology*, 47/4: 999-1014. <https://doi.org/10.1111/j.0031-0239.2004.00393.x>
- Desmarest, A.-G. 1817: Crustacés fossiles. In: Société de Naturalistes et d'Agriculteurs (ed.): Nouveau dictionnaire d'histoire naturelle, appliquée aux arts, à l'agriculture, à l'économie rurale et domestique, à la médecine, etc. 7. Déterville, Paris: 495-519.
- Desmarest, A.-G. 1822: Les crustacés proprement dits. In: Brongniart, A. & Desmarest, A.-G. (eds.): Histoire naturelle des crustacés fossiles sous les rapports zoologiques et géologiques. Levrault, Paris: 67-142.
- Devillez, J. & Charbonnier S. 2017: The genus *Eryma* Meyer, 1840 (Crustacea: Decapoda: Erymidae): new synonyms, systematic and stratigraphic implications. *Bulletin de la Société géologique de France*, 188/3: 1-10. <https://doi.org/10.1051/bsgf/2017178>
- Devillez, J. & Charbonnier, S. 2019: Review of the Early and Middle Jurassic erymid lobsters (Crustacea: Decapoda). *BSGF, Earth Sciences Bulletin*, 190/1: 6 <https://doi.org/10.1051/bsgf/2019005>
- Devillez, J. & Charbonnier, S. 2020: Review of the Late Jurassic erymid lobsters (Crustacea: Decapoda). *Geodiversitas*.
- Devillez, J., Charbonnier, S. & Pezy, J.-P. 2018: First Jurassic occurrence of *Enoploclytia* M'Coy, 1849 (Crustacea: Decapoda: Erymidae). *Annales de Paléontologie*, 104/2: 143-148. <https://doi.org/10.1016/j.annpal.2018.01.003>
- Dietl, G. & Schweigert, G. 2011: Im Reich der Meerengel - Fossilien aus dem Nusplinger Plattenkalk (2nd edition). Friedrich Pfeil, München: 144 p.
- Dumortier, E. 1867: Études paléontologiques sur les dépôts jurassiques du Bassin du Rhône, deuxième partie, Lias Inférieur. F. Savy, Paris: 252 p.
- Étallon, A. 1859: Description des crustacés fossiles de la Haute-Saône et du Haut-Jura. *Bulletin de la Société géologique de France*, 2/16: 169-204.
- Étallon, A. 1861: Notes sur les crustacés jurassiques du bassin du Jura. *Recueil agronomique, industriel et scientifique publié par la Société d'Agriculture de la Haute-Saône*, 9: 129-171.
- Feldmann, R.M. & Haggart, J.W. 2008: A new species of lobster (Astacidea, Erymidae) from the Smithers Formation (Middle Jurassic) of British Columbia, Canada. *Canadian Journal of Earth Sciences*, 44/12: 1791-1796. <https://doi.org/10.1139/E07-058>
- Fesefeldt, K. 1962: Schichtenfolge und Lagerung des oberen Weißjura zwischen Solnhofen und der Donau (südliche Frankenalb). *Erlanger geologische Abhandlungen*, 46: 1-80.
- Förster, R. 1966: Über die Erymiden, eine alte konservative Familie der mesozoischen Dekapoden. *Palaeontographica, Abteilung A*, 125/4-6: 61-175.
- Fraas, O. 1855: Beiträge zum obersten weissen Jura in Schwaben. *Jahreshefte des Vereins für vaterländische Naturkunde in Württemberg*, 11: 76-107.
- Frickhinger, K.A. 1994: Die Fossilien von Solnhofen. Goldschneck-Verlag, Korb: 336 p.
- Frickhinger, K.A. 1999: Die Fossilien von Solnhofen, 2nd volume. Goldschneck-Verlag, Korb: 190 p.
- Fürsich, F.T., Mäuser, M., Schneider, S. & Werner, W. 2007: The Wattendorf Plattenkalk (Upper Kimmeridgian) - a new conservation lagerstätte from the northern Franconian Alb, southern Germany. *Neues Jahrbuch für Geologie und Paläontologie Abhandlungen*, 245/1: 45-58. <https://doi.org/10.1127/0077-7749/2007/0245-0045>
- Haug, C., Haug, J.T., Waloszek, D., Maas, A., Frattigiani, R. & Liebau, S. 2009: New methods to document fossils from lithographic limestones of southern Germany and Lebanon. *Palaeontologia Electronica*, 12/3, 6T: 12 p. https://palaeo-electronica.org/2009_3/193/index.html
- Haug, J.T., Haug, C., Waloszek, D. & Schweigert, G. 2011: The importance of lithographic limestones for revealing ontogenies in fossil crustaceans. *Swiss Journal of Geosciences*, 104 (Supplement 1): 585-598. <https://doi.org/10.1007/s00015-010-0033-1>
- Hyžný, M., Schlögl, J., Charbonnier, S., Schweigert, G., Rulleau, L. & Gouttenoire, M. 2015: Intraspecific variation and taphonomy of a new erymid lobster (Crustacea: Decapoda) from the Middle Jurassic of Belmont (Beaujolais, France). *Geobios*, 48/5: 371-384. <https://doi.org/10.1016/j.geobios.2015.07.006>
- Krause, H.P. G. 1891: Die Decapoden des nord-deutschen Jura. *Zeitschrift der Deutschen Geologischen Gesellschaft*, 43: 171-225.
- Latreille, P.A. 1802-1803: Histoire naturelle, générale et particulière des Crustacés et des Insectes, 3. F. Dufart, Paris: 468 p.

- Leich, H. 1968: Nach Millionen Jahren ans Licht. Ott-Verlag, Thun & München: 180 p.
- M'Coy, F. 1849: On the classification of some British fossil Crustacea, with notices of new forms in the University collection at Cambridge. *Annals and Magazine of Natural History, including Zoology, Botany, and Geology*, 4/2: 330-335.
- Meyer, H. von 1861: [Briefliche Mittheilung an Prof. Bronn]. *Neues Jahrbuch für Mineralogie, Geognosie, Geologie und Petrefakten-Kunde*, 1861: 561, 678-679.
- Moret, L. 1946: *Eryma delphinensis* nouvelle espèce de Crustacé Décapode du Berriasien de Noyarey (Isère). *Travaux du Laboratoire de Géologie, Faculté des Sciences de Grenoble*, 25: 49-51.
- Münster, G. Graf zu. 1839: Decapoda Macroura. Abbildung und Beschreibung der fossilen langschwänzigen Krebse in den Kalkschiefern von Bayern. *Beiträge zur Petrefaktenkunde*, 2: 1-88.
- Niebuhr, B. & Pürner, T. 2014: Plattenkalk und Frankendolomit - Lithostratigraphie der Weißjura-Gruppe der Frankenalb (außer-alpiner Oberjura, Bayern). *Schriftenreihe der Deutschen Gesellschaft für Geowissenschaften*, 83: 5-71.
- Odin, G. P., Charbonnier, S., Devillez, J. & Schweigert, G. 2019: On new historical specimens of marine arthropods from the Solnhofen and Nusplingen Lithographic Limestones (Late Jurassic, Germany) housed at the Muséum national d'Histoire naturelle, Paris. *Geodiversitas*, 41/1: 643-662. <https://doi.org/10.5252/geodiversitas2019v41a17>
- Oppel, A. 1861: Die Arten der Gattungen *Eryma*, *Pseudastacus*, *Magila* und *Etallonia*. *Jahreshefte des Vereins für vaterländische Naturkunde in Württemberg*, 17: 355-361.
- Oppel, A. 1862: Ueber jurassische Crustaceen (Decapoda macrura). *Palaeontologische Mittheilungen aus dem Museum des königlich Bayerischen Staates*, 1: 1-120.
- Scherzinger, A. & Schweigert, G. 2003: Ein Profil in der Usseltal- und Rennertshofen-Formation der südlichen Frankenalb (Unter-Tithonium). *Zitteliana*, 43: 3-16. <https://doi.org/10.5282/ubm/epub.12120>
- Schlotheim, E. F. von. 1822: Beiträge zur näheren Bestimmung der versteinerten und fossilen Krebsarten. *Nachträge zur Petrefaktenkunde*. Beckersche Buchhandlung, Gotha: 17-37.
- Schweigert, G. 2001a: Dimorphismus bei Krebsen der Gattung *Cycleryon* (Decapoda, Eryonidae) aus dem Oberjura Süddeutschlands. *Stuttgarter Beiträge zur Naturkunde*, B305: 1-21.
- Schweigert, G. 2001b: Eine neue Art der Gattung *Antrimpos* Münster (Crustacea, Decapoda, Penaeidae) aus dem Oberjura Süddeutschlands. *Stuttgarter Beiträge zur Naturkunde*, B307: 1-33.
- Schweigert, G. 2006: Kalibrierung des Oberen Jura in der Stratigraphischen Tabelle von Deutschland 2002 mittels Faunenhorizonten. *Newsletters of Stratigraphy*, 41/1-3: 279-286. <https://doi.org/10.1127/0078-0421/2005/0041-0279>
- Schweigert, G. 2007: Ammonite biostratigraphy as a tool for dating Upper Jurassic lithographic limestones from South Germany - first results and open questions. *Neues Jahrbuch für Geologie und Paläontologie Abhandlungen*, 245/1: 117-125. <https://doi.org/10.1127/0077-7749/2007/0245-0117>
- Schweigert, G. 2013: A new record of the enigmatic lobster genus *Stenodactylina* Beurlen, 1928 (Crustacea: Decapoda: Erymidae) from the Middle Jurassic of south-western Germany. *Paläontologische Zeitschrift*, 87: 409-413. <https://doi.org/10.1007/s12542-013-0163-7>
- Schweigert, G. 2015a: Biostratigraphie der Plattenkalke der Südlichen Frankenalb. In: Arratia, G., Schultze, H.-P., Tischlinger, H. & Viohl, G. (eds.): *Solnhofen - Ein Fenster in die Jurazeit*. Verlag Friedrich Pfeil, München: 63-66.
- Schweigert, G. 2017: Die Garnele *Antrimpos undenarius* aus dem Nusplinger Plattenkalk (Ober-Kimmeridgium, Schwäbische Alb) und deren Rolle im dortigen Ökosystem. *Archaeopteryx*, 34: 1-7.
- Schweigert, G. & Dietl, G. 1999: Neubeschreibung von „*Eryon-longipes* O. Fraas“ (Crustacea, Decapoda, Eryonidea) aus dem Nusplinger Plattenkalk (Ober-Kimmeridgium, Schwäbische Alb). *Stuttgarter Beiträge zur Naturkunde*, B274: 1-19.
- Schweigert, G., Dietl, G. & Röper, M. 2000: Die Panzerkrebse der Familie Erymidae van Straelen (Crustacea, Decapoda) aus dem Nusplinger Plattenkalk (Ober-Kimmeridgium, Schwäbische Alb) im Vergleich mit fränkischen Vorkommen. *Stuttgarter Beiträge zur Naturkunde* B285: 1-25.
- Schweigert, G. & Garassino, A. 2003: New studies of decapod crustaceans from the Upper Jurassic lithographic limestones of southern Germany.

- Contributions to Zoology, 72/2: 173-179. <https://doi.org/10.1163/18759866-0720203022>
- Secretan, S. 1964: Les Crustacés décapodes du Jurassique supérieur et du Crétacé de Madagascar. Mémoires du Muséum national d'Histoire naturelle Paris, nouvelle série, C, Sciences de la Terre, 14: 1-226.
- Seilacher, A., Reif, W.-E. & Westphal, F. 1985: Sedimentological, ecological and temporal patterns of fossil Lagerstätten. Philosophical Transactions of the Royal Society of London, B311: 5-23.
- Stenzel, H.B. 1945: Decapod Crustacea from the Cretaceous of Texas. Texas University Publications, 4401: 401-476.
- Stevens, K., Mutterlose, J. & Schweigert, G. 2014: Belemnite ecology and the environment of the Nusplingen Plattenkalk (Late Jurassic, southern Germany): evidence from stable isotope data. Lethaia, 47/4: 512-523. <https://doi.org/10.1111/let.12076>
- Tischlinger, H. 2015: Arbeiten mit ultraviolettem Licht (UV). In: Arratia, G., Schultze, H.-P., Tischlinger, H. & Viohl, G. (eds.): Solnhofen - Ein Fenster in die Jurazeit. Friedrich Pfeil, München: 109-113.
- Van Straelen, V. 1925: Contribution à l'étude des crustacés décapodes de la période jurassique. Mémoires de la Classe des Sciences de l'Académie royale de la Belgique, 7: 1-462.
- Wright, T. 1881: On a new astacomorphous crustacean from the middle coral reef of Leckhampton Hill. Proceedings of the Cotteswold Naturalists' Field Club, 8: 56-59.
- Zeiss, A. 1968: Untersuchungen zur Paläontologie der Cephalopoden des Unter-Tithon der Südlichen Frankenalb. Abhandlungen der Bayerischen Akademie der Wissenschaften, mathematisch-naturwissenschaftliche Klasse, neue Folge, 132: 1-190.
- Zeiss, A. 1977: Jurassic stratigraphy of Franconia. Stuttgarter Beiträge zur Naturkunde, B31: 1-32.